Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

- 1-2. (Cancelled)
- 3. (Currently amended) A method for producing 3-hydroxypropionic acid comprising the steps of

providing in a fermenter a recombinant microorganism which carries a genetic construct which expresses the *dhaB* gene from *Klebsiella pneumoniae* and a gene for an aldehyde dehydrogenase, which are capable of catalyzing the production of 3-hydroxypropionic acid from glycerol;

providing a source of glycerol or glucose for the recombinant microorganism, and fermenting the microorganism under conditions which result in the accumulation of 3-hydroxypropionic acid <u>in solution in the fermenter</u>.

4. (Currently Amended) The A method of claim 3 for producing 3-hydroxypropionic acid comprising the steps of

providing in a fermenter a recombinant microorganism which carries a genetic construct which expresses the *dhaB* gene from *Klebsiella pneumoniae* and a gene for an aldehyde dehydrogenase, which are capable of catalyzing the production of 3-hydroxypropionic acid from glycerol;

providing a source of glycerol or glucose for the recombinant microorganism, and fermenting the microorganism under conditions which result in the accumulation of 3-hydroxypropionic acid wherein the gene for the aldehyde dehydrogenase is selected from the group consisting of ALDH4-ALDH2, ALD2ALD4, aldA and aldB.

5. (Currently Amended) The A method of claim 3 for producing 3-hydroxypropionic acid comprising the steps of

providing in a fermenter a recombinant microorganism which carries a genetic construct which expresses the *dhaB* gene from *Klebsiella pneumoniae* and a gene for an aldehyde dehydrogenase, which are capable of catalyzing the production of 3-hydroxypropionic acid from glycerol;

providing a source of glycerol or glucose for the recombinant microorganism, and fermenting the microorganism under conditions which result in the accumulation of 3-hydroxypropionic acid wherein the aldehyde dehydrogenase is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.

- 6. (Currently Amended) A recombinant *E. coli* host comprising in its inheritable genetic materials foreign genes encoding the *dhaB* gene from *Klebsiella pheumoniae* a glycerol dehydratase and an aldehyde dehydrogenase, such that the host is capable of producing 3-hydroxypropionic acid from glycerol.
- 7. (Original) A recombinant *E. coli* host comprising in its inheritable genetic materials the *dhaB* gene from *Klebsiella pheumoniae* and the *ald4* gene from *Saccharomycetes cervisiae*, such that the host is capable of producing 3-hydroxypropionic from glycerol.
- 8. (Currently Amended) A bacterial host comprising in its inheritable genetic material a genetic construction encoding for the expression of the dhaB gene from Klebsiella pheumoniae a glycerol dehydratase enzyme and an aldehyde dehydrogenase enzyme, such that the bacterial host is capable of converting glycerol to 3-hydroxypropionic acid.

9-10. (Cancelled)

- 11. (Currently Amended) The A bacterial host of claim 8 comprising in its inheritable genetic material a genetic construction encoding for the expression of a glycerol dehydratase enzyme and an aldehyde dehydrogenase enzyme, such that the bacterial host is capable of converting glycerol to 3-hydroxypropionic acid wherein the aldehyde dehydrogenase is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.
- 12. (Currently Amended) The A bacterial host of claim 8 comprising in its inheritable genetic material a genetic construction encoding for the expression of a glycerol dehydratase enzyme and an aldehyde dehydrogenase enzyme, such that the bacterial host is capable of converting glycerol to 3-hydroxypropionic acid wherein the gene for the aldehyde

dehydrogenase is selected from the group consisting of *ALDH4ALDH2*, *ALD2ALD4*, *aldA* and *aldB*.